

# Folding RNA/DNA hybrid duplexes

Ronny Lorenz<sup>1,\*</sup>, Ivo L Hofacker<sup>1,2</sup> and Stephan H Bernhart<sup>2,\*</sup>

<sup>1</sup>Institute for Theoretical Chemistry, University Vienna, Währinger Straße 17, 1090 Vienna, Austria

<sup>2</sup>Research group BCB, Faculty of Computer Science, University Vienna

<sup>3</sup>Department of Bioinformatics, University of Leipzig, Haertelstraße 16-18, 04109 Leipzig, Germany

Received on XXXXX; revised on XXXXX; accepted on XXXXX

Associate Editor: XXXXXXXX

## ABSTRACT

**Motivation:** While there are numerous programs that can predict RNA or DNA secondary structures, a program that predicts RNA/DNA hetero-dimers is still missing. The lack of easy to use tools for predicting their structure may be in part responsible for the small number of reports of biologically relevant RNA/DNA hetero-dimers.

**Results:** We present here an extension to the widely used ViennaRNA Package [6] for the prediction of the structure of RNA/DNA hetero-dimers.

**Availability:** <http://www.tbi.univie.ac.at/~ronny/RNA/vrna2.html>

**Contact:** [ronny@tbi.univie.ac.at](mailto:ronny@tbi.univie.ac.at)

## 1 INTRODUCTION

Nucleic acids have many important functions in biological systems, such as information carriers, catalysts and regulators. Both variants, DNA and RNA, can form complex structures by base pair interactions. The pattern of Watson-Crick or wobble base pairs a molecule forms is called the secondary structure. A main difference between RNA and DNA is due to their usage in biological systems: while DNA builds a dimer with its reverse complement almost all the time, RNA, being mostly single stranded, is more prone to fold back onto itself. However, if DNA is single stranded (e.g. during replication, transcription, repair or recombination), it will also form intramolecular base pairs. There are several examples where RNA and DNA interact via base pairing, generating a structure called R-loop. They protect CpG islands from being methylated [2]. The stability of DNA/mRNA hybrids is reported to have an influence on the copy number of repeats [8] and can impair transcription elongation, promoting transcription-associated recombination [5]. Moreover, during the intensive search for new functional transcripts rendered possible by next generation sequencing technology, at least 1000 novel lincRNAs were identified that may act as regulators of transcription, and one possibility to do this is via base pairing between RNA and DNA (see [3] for a recent review). As a final biological example, DNA/RNA hybrids play a role within the CRISPR pathway in bacteria and archaea [4]. On a technical side, RNA/DNA dimers are often used in micro-array or PCR experiments. The ability to computationally investigate the properties of the structures of interacting RNA and DNA molecules would help in the analysis of these dimers. While a number of programs exist that can predict RNA or DNA secondary structures

(for a review see e.g. [11]) or even their dimers, there is, to our knowledge, no program that also includes the possibility to predict the full secondary structure of RNA/DNA hetero-dimers. This may be due to the lack of compiled energy parameters, as there are only stacking energies available today. We introduce here a possibility to predict the secondary structure of such hetero-dimers within the widely used ViennaRNA Package.

## 2 APPROACH


### 2.1 Adaptation of the ViennaRNA Package

For the prediction of RNA/DNA hetero-dimers, three distinct energy parameter sets (RNA, DNA and mixed) are necessary. In order to keep the changes within the inner recursions of the algorithm minimal, we use 8 instead of four bases and 24 instead of six types of Watson-Crick or wobble base pairs for the energy computations. The look-up tables for the energy parameters grow accordingly. However, this does not significantly contribute to memory consumption. Finally, we demand a fixed order for the input sequences: RNA first, DNA last.

### 2.2 Energy parameters

Because we are only considering RNA/DNA hetero-dimers, not copolymers, we only need a subset of the energy parameters necessary for the full computations in the RNA or DNA case: we do not encounter stacking of a RNA on a DNA base, for example. In contrast, stacking parameters of RNA/DNA base pairs, multi-loop penalties and the interior loop parameters are needed. For stacked pairs, i.e. perfect helices, the parameters have been determined experimentally [7, 12]. Stacking parameters are the most important part of the energy model, as they account for the majority of RNA/DNA binding free energy. Unfortunately, for other mixed RNA-DNA loops no experimental data is available. As no high quality data set of RNA/DNA structures exists, parameters can not be trained, and the performance of different parameter sets can not be evaluated. The data we have is not sufficient to chose a function for the derivation of the energy parameters (supp. fig. 1), so we simply use the average of RNA and DNA  $E_{R/D} = (E_R + E_D)/2$  for all missing values. We are aware that this formula gives an at best crude approximation of the actual folding energies, but note that new parameters can easily be incorporated. We hope that the availability of prediction tools for RNA/DNA hetero-dimers will encourage experimentalists to provide the missing energy parameters.

\*to whom correspondence should be addressed



**Fig. 1.** Using hybrid instead of RNA parameters leads to very different structure prediction as summarized in a boxplot of the tree edit distances (A) for all interactions between an ncRNA and a putative target region at human chr. XX to YY (20000 nt). **B** to **D**: Predicted secondary structures for one example binding site (green: ncRNA), **B**: DNA parameters, **C**: RNA parameters, **D**: mixed parameters

### 2.3 Inclusion into the ViennaRNA Package

Our approach to just adapt the energy computations enabled us to easily include the RNA/DNA hetero-dimer support into all parts of the ViennaRNA Package that compute dimer structures, namely RNACoFold [1], RNAup [9] and RNAduplex [13]. RNA/DNA hetero-dimers are supported for both the minimum free energy as well as the partition function based calculations. Furthermore, all programs can read in user supplied energy parameters. In particular, this allows to make immediate use of any newly determined parameters for mixed RNA-DNA loops, or enables scientists to easily evaluate different energy models as soon as sufficient RNA/DNA structure data is available.

## 3 RESULTS

In the absence of programs that can deal with RNA/DNA dimers, researchers have resorted to simply treating both parts of the dimer as RNA. To test the effect of applying the right (DNA or RNA) parameters to the single molecules and using explicit RNA/DNA interactions, we chose a long non-coding RNA that is suspected to have a DNA target sequence, and one of its putative targets. In order to simulate the size of the transcription bubble, we chose a sliding window approach with DNA pieces of length 50, step size 25 for our computations. Even though the ncRNA (1500nt) is much longer than the pieces of DNA bound to it, we sometimes observe a dramatic effect on the structure of the hybrid molecule. The tree edit distance of the hybrid structures predicted with RNA parameters and with our hybrid approach rises up to 900, and the structure of the binding site can vary greatly (Fig 1).

Another example for the usefulness of the mixed parameters is the strand dependency of R-loop formation [10]. A  $r(GAA)_n$  trinucleotide repeat can form R-loops while its reverse complement  $r(UUC)_n$  cannot. The non-symmetric mixed parameters can explain that: addition of another trinucleotide will give  $-4.1$  kcal/mol for  $r(GAA)$ , but only  $-2.6$  kcal/mol for  $r(UUC)$ . In contrast to RNA or DNA parameters, the mixed parameters can also explain the changes in R-loop formation of other trinucleotides investigated (see supp. fig. 2).

## ACKNOWLEDGMENT

*Funding:* This work was funded, in part, by the austrian “SFB 43 RNA regulation of the transcriptome” and the austrian GEN-AU project “Bioinformatics Integration Network III”.

## REFERENCES

- [1] S H Bernhart, H Tafer, U Mückstein, C Flamm, P F Stadler, and I L Hofacker. Partition function and base pairing probabilities of RNA heterodimers. *Algorithms Mol Biol*, 1(1):3–3, 2006.
- [2] P A Ginno, P L Lott, H C Christensen, I Korf, and F Chédin. R-loop formation is a distinctive characteristic of unmethylated human CpG island promoters. *Mol Cell*, Feb 2012.
- [3] M Guttman and J L Rinn. Modular regulatory principles of large non-coding RNAs. *Nature*, 482(7385):339–346, Feb 2012.
- [4] J A Howard, S Delmas, I Ivani-Bae, and E L Bolt. Helicase dissociation and annealing of RNA-DNA hybrids by escherichia coli cas3 protein. *Biochem J*, 439(1):85–95, Oct 2011.
- [5] P Huertas and A Aguilera. Cotranscriptionally formed DNA:RNA hybrids mediate transcription elongation impairment and transcription-associated recombination. *Mol Cell*, 12(3):711–721, Sep 2003.
- [6] R Lorenz, S H Bernhart, C Höner Zu Siederdisen, H Tafer, C Flamm, P F Stadler, and I L Hofacker. ViennaRNA package 2.0. *Algorithms Mol Biol*, 6(1):26–26, Nov 2011.
- [7] F H Martin and I Tinoco. DNA-RNA hybrid duplexes containing oligo(da:ru) sequences are exceptionally unstable and may facilitate termination of transcription. *Nucleic Acids Res*, 8(10):2295–2299, May 1980.
- [8] E I McIvor, U Polak, and M Napierala. New insights into repeat instability: role of RNADNA hybrids. *RNA Biol*, 7(5):551–558, Sep-Oct 2010.
- [9] U. Mückstein, H. Tafer, S. H. Bernhart, M. Hernandez-Rosales, J. Vogel, P. F. Stadler, and I. L. Hofacker. Translational control by RNA-RNA interaction: Improved computation of RNA-RNA binding thermodynamics. In M. Elloumi, J. Küng, M. Linial, R. Murphy, K. Schneider, and C. Toma, editors, *Bioinformatics Research and Development*, volume 13 of *Communications in Computer and Information Science*, pages 114–127. Springer, 2008.
- [10] K Reddy, M Tam, R P Bowater, M Barber, M Tomlinson, K Nichol Edamura, Y H Wang, and C E Pearson. Determinants of r-loop formation at convergent bidirectionally transcribed trinucleotide repeats. *Nucleic Acids Res*, 39(5):1749–1762, Mar 2011.
- [11] J. Reeder, M. Höchsmann, M. Rehmsmeier, B. Voss, and R. Giegerich. Beyond Mfold: recent advances in RNA bioinformatics. *Journal of Biotechnology*, 124(1):41–55, 2006.
- [12] N Sugimoto, S Nakano, M Katoh, A Matsumura, H Nakamura, T Ohmichi, M Yoneyama, and M Sasaki. Thermodynamic parameters to predict stability of RNA/DNA hybrid duplexes. *Biochemistry*, 34(35):11211–11216, Sep 1995.
- [13] H. Tafer and I. L. Hofacker. RNAplex: a fast tool for RNA-RNA interaction search. *Bioinformatics*, 24(22):2657–2663, 2008.